

Problem: We want a forward difference expression for the **first** derivative that is accurate to $\mathcal{O}(h^2)$.

Starting Point: $f(x+h) = f(x) + hf'(x) + \frac{h^2}{2!} f''(x) + \frac{h^3}{3!} f'''(x) + \frac{h^4}{4!} f^{iv}(x) + \dots$ [1]

Similarly, $f(x+2h) = f(x) + 2hf'(x) + \frac{(2h)^2}{2!} f''(x) + \frac{(2h)^3}{3!} f'''(x) + \frac{(2h)^4}{4!} f^{iv}(x) + \dots$ [2]

Solving [1] for f': $f'(x) = \frac{f(x+h) - f(x)}{h} - \frac{h}{2} f''(x) - \frac{h^2}{6} f'''(x) - \frac{h^3}{24} f^{iv}(x) + \dots$ [3]

Algebraically performing EQ [2] - 2 × EQ[1] results in:

$$f(x+2h) - 2f(x+h) = [f(x) - 2f(x)] + [2hf'(x) - 2hf'(x)] + \left[\frac{(2h)^2}{2!} f''(x) - 2 \frac{h^2}{2!} f''(x) \right] + \left[\frac{(2h)^3}{3!} f'''(x) - 2 \frac{h^3}{3!} f'''(x) \right] + \dots$$

This simplifies to:

$$f(x+2h) - 2f(x+h) = [-f(x)] + [0] + [h^2 f''(x)] + [\mathcal{O}(h)^3] = -f(x) + h^2 f''(x) + \mathcal{O}(h^3)$$

Rearranging results in: $f''(x) = \frac{f(x+2h) - 2f(x+h) + f(x)}{h^2} + \mathcal{O}(h^1)$ [4]

Next, we substitute our EQ[4] into EQ[3]:

$$f'(x) = \frac{f(x+h) - f(x)}{h} - \frac{h}{2} \left[\frac{f(x+2h) - 2f(x+h) + f(x)}{h^2} \right] + \mathcal{O}(h^2)$$
 [5]

Expanding and distributing:

$$f'(x) = \frac{f(x+h) - f(x) + \left[-\frac{1}{2} f(x+2h) + f(x+h) - \frac{1}{2} f(x) \right]}{h} + \mathcal{O}(h^2) + \mathcal{O}(h^2)$$

One more rearrangement:

$$f'(x) = \frac{-f(x+2h) + 4f(x+h) - 3f(x)}{2h} + \mathcal{O}(h^2)$$

Rewriting with subscript notation:

$f'_i = \frac{-f_{i+2} + 4f_{i+1} - 3f_i}{2h} + \mathcal{O}(h^2)$
