

Strongly Uncontrollable Network Topologies

Cesar O. Aguilar 

Abstract—In this article, we present a class of network topologies under which the Laplacian consensus dynamics exhibit undesirable controllability properties under a broadcast control signal. Specifically, the networks we characterize are uncontrollable for any subset of the nodes chosen as control inputs and that emit a common control signal. We provide a sufficient condition for a network to contain this strong uncontrollability property and describe network perturbations that leave the uncontrollability property invariant. As a by-product, we identify nontrivial network topologies that require the control of approximately half the nodes in the network as a necessary condition for controllability.

Index Terms—Controllability, consensus dynamics, graph theory, Laplacian matrix, multiagent systems.

I. INTRODUCTION

CONTROLLABILITY of networked multiagent systems is an ongoing topic of research in the control systems community due to the proliferation of technologies associated with large-scale network models, see for instance [1]–[13] and references therein. A primary goal of the ongoing research has been to identify graph-theoretic structures that are possible obstructions to controlling a networked control system. To that end, the lack of controllability has been primarily attributed to the existence of so-called *equitable partitions* of the vertex set [2], [3], [6], [8], [10]. We note that a special case of these equitable partitions is the presence of structural symmetries in the associated network model [2]. Roughly speaking, the existence of an equitable partition of the nodes of a network induce an invariant subspace for the uncontrolled dynamics and, thus, if the control nodes are chosen to preserve the invariant subspace, then uncontrollability ensues. A closely related line of research is the so-called *minimal controllability problem* which is concerned with the scenario of controlling a large-scale multiagent system with the fewest possible number of control nodes [14]–[18]. Although it has been shown that solving minimal controllability problems is computationally intractable for generic systems (unless $P = NP$), heuristic algorithms are known that produce approximate solutions [14], [16]–[18]. On the other hand, in the case of structured systems, the minimal controllability problem can be solved in polynomial time [15], [19].

Manuscript received June 7, 2019; revised August 28, 2019 and August 29, 2019; accepted October 29, 2019. Date of publication November 6, 2019; date of current version June 12, 2020. This work was supported by the National Science Foundation under Grant ECCS-1700578. Recommended by Associate Editor Dr. Dragan Nesic.

The author is with the Department of Mathematics, State University of New York, Geneseo, NY 14454 USA (e-mail: aguilar@geneseo.edu).
Digital Object Identifier 10.1109/TCNS.2019.2951665

TABLE I

$s(n)$ IS THE NUMBER OF CONNECTED GRAPHS ON n VERTICES SUCH THAT $L = L(G)$ HAS DISTINCT EIGENVALUES AND (L, b) IS UNCONTROLLABLE FOR every $b \in \{0, 1\}^n$

n	8	9	10	11	12
$s(n)$	10	12	91	232	1749

In this article, we are motivated by the following question: What structural properties present in a network result in uncontrollable dynamics for any choice of control nodes emitting a common control signal? Specifically, we focus on networked multiagent systems undergoing the Laplacian consensus dynamics and describe network topologies that are uncontrollable under a broadcast control for any choice of leader nodes. Such networks were introduced in [9] and were called *strongly uncontrollable graphs*. To be more precise about the issue at hand, let G be a simple n -vertex graph with Laplacian matrix $L = L(G) = A(G) - D(G)$, where $A(G)$ is the adjacency matrix and $D(G)$ is the diagonal degree matrix of G , and let b be a binary vector. A trivial necessary condition for the pair (L, b) to be controllable is that the eigenvalues of L are all distinct [20, p. 95]. Aguilar and Ghahesifard [9], however, provide examples of graphs for which L has *distinct* eigenvalues but (L, b) is uncontrollable for every choice of binary vector b and such graphs were called strongly uncontrollable graphs.¹ A similar definition of *strong uncontrollability* can be made using the adjacency matrix $A(G)$, the signless Laplacian matrix $Q(G) = A(G) + D(G)$, or some other graph matrix relevant to the network model in consideration. In the case of the Laplacian matrix, we have performed an exhaustive numerical search revealing that strongly uncontrollable graphs do not appear until the number of nodes is $n = 8$, that is, no connected graph is strongly uncontrollable for $n \leq 7$ in the case of L . Our numerical search produced an enumeration of strongly uncontrollable graphs for $8 \leq n \leq 12$ and the results are shown in Table I. Interestingly, for the case of the adjacency matrix, our numerical search revealed that no connected graph for $n \leq 10$ is strongly uncontrollable and, based on our results in this article, we conjecture that strongly uncontrollable graphs using $A(G)$ do not exist.

The main contribution of this article is the characterization of a class of network topologies that result in a strongly uncontrollable multiagent system undergoing the Laplacian consensus dynamics. The identification of these topologies could provide a

¹In [9, Theor. 4.1], there is also a construction of a class of graphs such that (L, b) is uncontrollable for every choice of binary vector b but such graphs contain *repeated* eigenvalues.

test bed to narrow the gap between known sufficient conditions and necessary conditions for network controllability. As a by-product of our results, we identify a class of nontrivial network topologies that require the control of approximately half of the nodes for any chance of controllability. The topologies we study are of interest since they contain many “local symmetries” that are similar to the symmetries found in large-scale real-world complex networks [21]. The discovered network topologies contain two main structural ingredients, namely, the presence of many so-called *twin* nodes and certain equitable partitions. Twin vertices induce so-called Faria eigenvectors [22] of the Laplacian matrix L and, through the use of the classical Popov-Belevitch-Hautus (PBH) test for controllability, result in the uncontrollability of (L, b) for many choices of b but, in general, are not enough for strong uncontrollability. The second main ingredient is the presence of certain equitable partitions [23] which, together with the presence of twin nodes, result in a strongly uncontrollable graph. Graph vertex partitions are becoming a standard tool used to study control-theoretic properties in multiagent systems, see, for instance, [2], [3], [6], [24], [25] and references therein. Graph vertex partitions also take an important role in the study of synchrony and pattern formation in coupled cell networks [26], [27]. In addition to our main result, and motivated by the recent research activity with structural controllability, we identify graph perturbations which preserve the strong uncontrollability property.

This article is organized as follows. In Section II, we introduce our notation, review the connection between equitable partitions of a graph and controllability, and present some technical results. In Section III, we introduce twin graphs, which are nontrivial network topologies requiring that approximately half of the nodes be controlled. In Section IV, we present our main result of the article, namely, a sufficient condition for a networked multiagent system undergoing the Laplacian consensus dynamics to be uncontrollable for any choice of leader nodes under a broadcast control signal. In Section V, we identify three vertex addition operations that leave the strong uncontrollability property invariant. We end the article with a Conclusion.

II. PRELIMINARIES

In this section, we introduce our notation, review the notion of equitable partitions and their role in network controllability, and present some technical results.

A. Notation

The all ones vector in \mathbb{R}^n is denoted by $e = (1, 1, \dots, 1)$ and the context will make it clear the value of n . For $u \in \{1, 2, \dots, n\}$, we denote by e_u the unit standard basis vector in \mathbb{R}^n with a nonzero entry at u . We say that a square matrix M has a *simple spectrum* if every eigenvalue of M has algebraic multiplicity one. The column/range space of M will be denoted by $\text{range}(M)$. We equip \mathbb{R}^n with the standard Euclidean inner product $\langle x, y \rangle = x^T y$ and denote by $\Omega^\perp = \{x \in \mathbb{R}^n \mid \langle x, y \rangle = 0, \forall y \in \Omega\}$ the orthogonal complement of a set $\Omega \subseteq \mathbb{R}^n$. A subspace $W \subset \mathbb{R}^n$ is said to be *M-invariant* if $w \in W$ implies that $Mw \in W$. For each positive

integer n , we let $\{0, 1\}^n$ denote the set of binary vectors of length n . Finally, we let $\mathbb{N}_0 = \{0, 1, 2, \dots\}$.

By a *weighted digraph*, we mean a triple $G = (V, E, \phi)$ where V is the set of vertices, $E \subset V \times V$ is the set of directed edges (or arcs), and $\phi : E \rightarrow \mathbb{N}_0$ is the weight function on the arcs with the property that $\phi(u, v) \neq 0$ if and only if $(u, v) \in E$. We do not have a need for loops in a graph and, thus, $(u, u) \notin E$ for all $u \in V$. If $(u, v) \in E$ if and only if $(v, u) \in E$ and $\phi \equiv 1$, then we call G a *simple graph* and instead use the usual notation $G = (V, E)$. The context will make it clear as to whether G is a simple graph or a weighted digraph.

Let $G = (V, E, \phi)$ be a weighted digraph. The *neighborhood* of $u \in V$ is the set $N(u) = \{v \in V \mid (u, v) \in E\}$ and the *degree* of u is $\deg(u) = \sum_{v \in V} \phi(u, v)$. For a simple graph G , $\deg(u) = |N(u)|$ for all $u \in V$, i.e., the number of vertices in $N(u)$. More generally, for a subset $C \subseteq V$, we define the degree of u in C by

$$\deg(u, C) := \sum_{v \in C} \phi(u, v).$$

If G is a simple graph, then $\deg(u, C) = |N(u) \cap C|$. If we label the vertices as $V = \{v_1, v_2, \dots, v_n\}$, the adjacency matrix $A = A(G)$ of G has entries $A_{i,j} = \phi(v_i, v_j)$. The degree matrix of G is the diagonal matrix $D = D(G)$ whose i th diagonal element is $\deg(v_i)$, and the Laplacian matrix of G is $L(G) = D - A$.

Finally, we recall that a linear single-input time-invariant system $\dot{x} = Ax + bu$ on \mathbb{R}^n is controllable if and only if the smallest A -invariant subspace containing b is all of \mathbb{R}^n , that is, $\text{span}\{b, Ab, \dots, A^{n-1}b\} = \mathbb{R}^n$. A well-known characterization of controllability is the PBH eigenvector test, which states that (A, b) is controllable if and only if $\xi^T b \neq 0$ for every eigenvector ξ of A^T .

B. Almost Equitable Partitions

Let G be a weighted digraph with vertex set $V = \{v_1, v_2, \dots, v_n\}$. The *characteristic vector* of a subset $C \subset V$ is the binary vector $\xi \in \{0, 1\}^n$ such that $\xi_i = 1$ if and only if $v_i \in C$. Let $\pi = \{C_1, C_2, \dots, C_k\}$ be a set partition of V , that is, $C_i \cap C_j = \emptyset$ if $i \neq j$ and $\bigcup_{i=1}^k C_i = V$. The subsets C_i will be called *cells* of π . The *characteristic matrix* of π is the $n \times k$ matrix P_π whose i th column is the characteristic vector of C_i , for $i = 1, \dots, k$. When no confusion arises, we denote P_π simply by P . We say that π is an *almost equitable partition* (AEP) of G if for every distinct ordered pair of cells (C_i, C_j) it holds that $\deg(u, C_j) = \deg(v, C_j)$ for every $u, v \in C_i$. In this case, we define $\deg(C_i, C_j) := \deg(u, C_j)$ for some (and hence all) $u \in C_i$. We note that, in general, $\deg(C_i, C_j) \neq \deg(C_j, C_i)$. If π is an AEP of G , the *quotient graph* G_π of G with respect to π is the weighted digraph with vertex set $V(G_\pi) = \pi$ and arcs $(C_i, C_j) \in E(G_\pi)$ if and only if $\deg(C_i, C_j) \neq 0$ with arc weight $\phi(C_i, C_j) = \deg(C_i, C_j)$. We denote the adjacency matrix of G_π by $A_\pi = A(G_\pi)$ and its Laplacian matrix by $L_\pi = L(G_\pi)$, and we note that A_π and L_π are generally nonsymmetric matrices. An AEP $\pi = \{C_1, \dots, C_k\}$ of G is called an *equitable partition* if in addition $\deg(u, C_i) = \deg(v, C_i)$ for all $u, v \in C_i$ and all $i = 1, \dots, k$. Hence, the extra




Fig. 1. Example of a graph G with an AEP π and the induced quotient graph G_π .

requirement imposed on an equitable partition is that vertices in any given cell C_i have equal degree *within* C_i . To make this paper as self-contained as possible, below we provide an example of the previous notions.

Example II.1: Consider the simple graph G shown in Fig. 1 with $n = 11$ vertices. Consider the vertex partition $\pi = \{C_1, C_2, C_3, C_4\}$ where $C_1 = \{v_1, v_2, v_3, v_4\}$, $C_2 = \{v_5, v_6\}$, $C_3 = \{v_7, v_8, v_9, v_{10}\}$, and $C_4 = \{v_{11}\}$. The edges within each cell are displayed as dashed lines. The reader is invited to verify that for any pair of distinct cells (C_i, C_j) it holds that $\deg(v, C_j) = \deg(u, C_j)$ for every $u, v \in C_i$. Notice that, for instance, $\deg(v_1, C_1) = 1$ while $\deg(v_3, C_1) = 2$, and, thus, π is an AEP of G but not an equitable partition. The characteristic matrix P of π is the $n \times k = 11 \times 4$ matrix

$$P^T = \begin{bmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

The quotient graph G_π is also shown in Fig. 1 and the adjacency and Laplacian matrix of G_π are

$$A_\pi = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 2 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 2 & 4 & 0 \end{bmatrix}, \quad L_\pi = \begin{bmatrix} 1 & -1 & 0 & 0 \\ -2 & 5 & -2 & -1 \\ 0 & -1 & 2 & -1 \\ 0 & -2 & -4 & 6 \end{bmatrix}.$$

We note that A_π and L_π are nonsymmetric matrices. ■

The relationship between AEPs and invariant subspaces of the Laplacian matrix L is the following.

Theorem II.1 (see[23]): Let G be a weighted digraph and let $\pi = \{C_1, C_2, \dots, C_k\}$ be a partition of $V(G)$ with characteristic matrix P . Then, π is an AEP of G if and only if $\text{range}(P)$ is L -invariant. In this case, the Laplacian matrix of the quotient graph G_π is $L_\pi = (P^T P)^{-1} P^T L P$.

Remark II.1: We make the following important observations as a consequence of Theorem II.1, which explains the connection between AEPs and obstructions to controllability. Let $G = (V, E)$ be a simple n -vertex graph, let $\pi = \{C_1, \dots, C_k\}$ be an AEP of G , and let P be the characteristic matrix of π . Since the subspace $\text{range}(P) \subset \mathbb{R}^n$ is L -invariant, there exists a basis for $\text{range}(P)$ consisting of eigenvectors of L . Moreover, since L is symmetric, then $\text{range}(P)^\perp = \ker(P^T)$ is also L -invariant and, thus, $\ker(P^T) \subset \mathbb{R}^n$ also has a basis of eigenvectors of L .

Now, by definition of P , $x \in \text{range}(P)$ if and only if the components of x are constant on each cell of π , that is, $\forall v_i, v_j \in C_\ell$, we have $x_i = x_j$ and this holds for all $\ell = 1, \dots, k$. Similarly, $x \in \ker(P^T)$ if and only if the components of x sum to zero on the cells of π , that is, $\sum_{v_j \in C_\ell} x_j = 0$ for each $\ell = 1, \dots, k$. Hence, if π is an AEP of G and L is a symmetric matrix, then the eigenvectors of L can be divided into two classes: one class is contained in $\text{range}(P)$ and are characterized by having a constant value on each cell of π , and the second class is contained in $\ker(P^T)$ and are characterized by summing to zero on each cell of π . Hence, if $b \in \{0, 1\}^n$ is constant on the cells of π , that is $b = P\bar{b}$ for some $\bar{b} \in \{0, 1\}^k$, then b is clearly orthogonal to the eigenvectors of L contained in $\ker(P^T)$ and, therefore, by the PBH test (L, b) is uncontrollable. We illustrate our remarks with our running example.

Example II.2: Consider again the simple graph G shown in Fig. 1, with AEP $\pi = \{C_1, C_2, C_3, C_4\}$ where $C_1 = \{v_1, v_2, v_3, v_4\}$, $C_2 = \{v_5, v_6\}$, $C_3 = \{v_7, v_8, v_9, v_{10}\}$, and $C_4 = \{v_{11}\}$. Hence, $x \in \text{range}(P_\pi)$ if and only if $x = (\alpha, \alpha, \alpha, \alpha, \beta, \beta, \gamma, \gamma, \gamma, \gamma, \delta)$ for some scalars $\alpha, \beta, \gamma, \delta \in \mathbb{R}$. Since $\text{rank}(P_\pi) = k = 4$, there are 4 linearly independent eigenvectors of L of the form of x above. One such vector is of course the all ones vector $e \in \mathbb{R}^{11}$ in which case $\alpha = \beta = \gamma = \delta = 1$. In general, since an eigenvector x of the form above (with eigenvalue $\lambda \neq 0$) is orthogonal to e , we must have $4\alpha + 2\beta + 4\gamma + \delta = 0$. On the other hand, $\ker(P^T)$ is a seven-dimensional subspace of \mathbb{R}^{11} and it can be verified that $\ker(P^T)$ has a basis consisting of eigenvectors of L . Since $C_4 = \{v_{11}\}$ is a singleton cell, all eigenvectors of L in $\ker(P^T)$ have a zero in the last entry. It follows that if, for example, $b = (0, 0, 0, 0, 1, 1, 0, 0, 0, 0, 1) \in \text{range}(P)$, then b is orthogonal to every eigenvector of L in $\ker(P^T)$ and, thus, (L, b) is uncontrollable by the PBH test.

There is a well-known relationship between the eigenvalues/eigenvectors of L and L_π , namely that (y, λ) is an eigenvector/eigenvalue pair of L_π if and only if (Py, λ) is an eigenvector/eigenvalue pair of L , where P is the characteristic matrix of π [23]. Now, it is possible that the quotient graph G_π itself contains an AEP and in this case an AEP of G_π induces an AEP of G in the following way. Let $\pi = \{C_1, \dots, C_k\}$ be a partition of V and let $\rho = \{S_1, \dots, S_m\}$ be a partition of π . We define the ρ -merge of π as the partition $\pi_\rho = \{\bar{C}_1, \bar{C}_2, \dots, \bar{C}_m\}$ of V where, for $j = 1, 2, \dots, m$, we have

$$\bar{C}_j = \bigcup_{C_i \in S_j} C_i.$$

Roughly speaking, the ρ -merge of π is simply obtained by “flattening out” each cell S_j of ρ . For example, if $\pi = \{C_1, C_2, C_3, C_4\} = \{\{1, 2, 3\}, \{4, 5\}, \{6, 7\}, \{8\}\}$ and $\rho = \{S_1, S_2\} = \{\{C_1, C_4\}, \{C_2, C_3\}\}$, then the ρ -merge of π is $\pi_\rho = \{\{1, 2, 3, 8\}, \{4, 5, 6, 7\}\}$. Notice that the partition π is *finer* than π_ρ , i.e., every cell in π is a subset of a cell of π_ρ . Now, if π is an AEP of G and ρ is an AEP of G_π , then it is known that the ρ -merge of π is an AEP of G [10, Prop. 1].

If ρ is an AEP of G_π , then, by Theorem II.1, $\text{range}(P_\rho)$ is L_π -invariant but it does not generally hold that the orthogonal complement $\text{range}(P_\rho)^\perp$ is L_π -invariant since L_π is not

generally a symmetric matrix. Hence, the discussion in Remark II.1 regarding the eigenvector structure of L does not generally hold for L_π , i.e., the eigenvectors of L_π do not generally split into those that are constant on the cells of ρ and those that sum to zero on the cells of ρ . There is, however, a case in which the eigenvectors of L_π do split into those in $\text{range}(P_\rho)$ and $\text{range}(P_\rho)^\perp$, and this case is present in the network topologies that we characterize. With this in mind, we introduce the following notion.

Definition II.1: Let $\pi = \{C_1, C_2, \dots, C_k\}$ be a partition of V . A partition $\rho = \{S_1, S_2, \dots, S_m\}$ of π is said to be π -regular if each cell $S_j \in \rho$ consists of cells of π of the same cardinality.

In other words, if $S_j = \{C_{j,1}, C_{j,2}, \dots, C_{j,m_j}\}$, then ρ is π -regular if $|C_{j,1}| = |C_{j,2}| = \dots = |C_{j,m_j}|$, and this holds for all $j = 1, 2, \dots, m$. As an example, if $V = \{1, 2, \dots, 12\}$ and $\pi = \{C_1, C_2, \dots, C_6\}$ where $C_1 = \{1, 2, 3\}$, $C_2 = \{4, 5\}$, $C_3 = \{6\}$, and $C_4 = \{7, 8\}$, $C_5 = \{9\}$, $C_6 = \{10, 11, 12\}$, then $\rho = \{\{C_1, C_6\}, \{C_2, C_4\}, \{C_3, C_5\}\}$ is π -regular since $|C_1| = |C_6|$, $|C_2| = |C_4|$, and $|C_3| = |C_5|$. The relationship between π -regularity of ρ and the L_π -invariance of $\text{range}(P_\rho)^\perp$ is then given in the following lemma whose proof can be found in Appendix A.

Lemma II.1: Let G be a simple graph. Let π be an AEP of G and let ρ be an AEP of the quotient graph G_π . If ρ is π -regular, then $\text{range}(P_\rho)^\perp = \ker(P_\rho^T)$ is L_π -invariant.

The upshot of Lemma II.1 is that even though L_π may not be a symmetric matrix, the discussion in Remark II.1 regarding the splitting structure of the eigenvectors of L is also applicable to L_π provided ρ is a π -regular AEP of G_π . Hence, if the conditions of Lemma II.1 are satisfied, then the eigenvectors of L_π can be partitioned into two classes, those contained in $\text{range}(P_\rho)$ and the others contained in $\ker(P_\rho^T)$.

III. TWIN GRAPHS

One of the main structural properties possessed by the network topologies that we characterize is the existence of many *twin vertices*. The vertices $u, v \in V(G)$ are *twins* if $N(u) \setminus \{v\} = N(v) \setminus \{u\}$. We note that it is possible for u to be twins with multiple vertices, that is, that $\{u, v\}$ and $\{u, w\}$ are twins with $v \neq w$. In this case, either $\{u, v, w\}$ are all mutually adjacent or nonadjacent. The existence of twin vertices induces an equitable partition as follows. Recall that a permutation $\sigma : V \rightarrow V$ is an *automorphism*, or *symmetry*, of the graph $G = (V, E)$ if $\{u, v\} \in E$ if and only if $\{\sigma(u), \sigma(v)\} \in E$ for all $u, v \in V$. We denote by $\text{Aut}(G)$ the group of automorphisms of G . It is clear that if u and v are twin vertices, then the permutation $\sigma : V \rightarrow V$ that transposes u and v and fixes all other vertices is an automorphism of G . Moreover, if without loss of generality $u = v_1$ and $v = v_2$, then the partition $\pi = \{\{v_1, v_2\}, \{v_3\}, \{v_4\}, \dots, \{v_n\}\}$ is an equitable partition of G . The following summarizes the relationship between twin vertices and eigenvectors/eigenvalues of L (see [22, pg. 46] for a proof).

Proposition III.1 (see[22]): Let $G = (V, E)$ be a simple graph with Laplacian matrix L , let $u, v \in V$, and let $\sigma : V \rightarrow V$ be the permutation that transposes u and v and fixes all other

vertices. Then, σ is an automorphism of G if and only if $x = e_u - e_v$ is an eigenvector of L . In this case, L has corresponding integer eigenvalue $\lambda = \deg(u) + A_{u,v}$.

An eigenvector of L of the form $x = e_u - e_v$ is known as a Faria eigenvector [22]. Clearly, a graph containing the Faria eigenvector $x = e_u - e_v$ will result in the uncontrollability of (L, b) for all binary vectors b whose entries are equal on u and v since then $\langle x, b \rangle = b_u - b_v = 0$. Note that clearly $x = e_u - e_v \in \ker(P_\pi^T)$ where $\pi = \{\{u, v\}, \{v_3\}, \{v_4\}, \dots, \{v_n\}\}$.

We now obtain an upper bound on the number of twin vertices in a graph whose Laplacian matrix has simple eigenvalues. To that end, we first recall that the *order* of a permutation $\sigma : V \rightarrow V$ is the smallest integer k such that $\sigma^k := \sigma \circ \sigma \circ \dots \circ \sigma = \text{id}$, where in the composition σ appears k -times.

Lemma III.1: Suppose that G is a simple graph on n vertices and assume that L has simple spectrum. The following hold:

- 1) If $\{u, v\}$ are twin vertices then $\deg(u) = \deg(v) < n - 1$.
- 2) If n is even then the maximal number of twins is $t = \frac{n}{2} - 1$, and this bound is sharp.
- 3) If n is odd then the maximal number of twins is $t = \frac{n-1}{2}$, and this bound is sharp.

Proof: To prove 1), suppose that $\deg(u) = \deg(v) = n - 1$ for a twin pair $\{u, v\} \subset V$. Then, by Proposition III.1, $\lambda = n$ is an eigenvalue of L with Faria eigenvector $x = e_u - e_v$. On the other hand, it is straightforward to verify that $\tilde{x} = -e + ne_u$ is also an eigenvector of L affording the eigenvalue $\lambda = n$. Thus, $\lambda = n$ has algebraic multiplicity at least 2, which contradicts the assumption that L has simple spectrum. Hence, we must have $\deg(u) = \deg(v) < n - 1$.

To prove 2), we first recall that if G has an automorphism of order $k \geq 3$, then L has a repeated eigenvalue [28, pg. 45]. Now, if G has more than $\frac{n}{2}$ twin pairs of vertices, then by the pigeon-hole principle, there are at least two sets of twin vertices of the form $\{u, v\}$ and $\{u, w\}$ with $v \neq w$. Then, the cyclic automorphism $\sigma(u) = v, \sigma(v) = w$, and $\sigma(w) = u$ (and all other vertices held fixed) clearly has order three and consequently L does not have a simple spectrum. Suppose now then that G has exactly $\frac{n}{2}$ twins and that L has a simple spectrum. We claim that every eigenvalue induced by a twin is even. To see this, if $\{u, v\}$ are twins, then $\deg(u) = 2q_u + A_{u,v}$ where q_u is the number of twins u (and hence v) is adjacent to. By Proposition III.1, the eigenvalue induced by $\{u, v\}$ is $\lambda = \deg(u) + A_{u,v} = 2(q_u + A_{u,v})$, which is even, and proves the claim. Since L has simple spectrum and all the eigenvalues induced by the $\frac{n}{2}$ twins are even, it follows that n is an eigenvalue of L . Thus, there exists a twin pair each of which has degree $n - 1$, which contradicts part 1). Hence, this proves that no simple graph with Laplacian simple eigenvalues has $\frac{n}{2}$ or more twin vertices. To prove that $\frac{n}{2} - 1$ is a sharp bound for the number of twins, it may be verified that the graph on $n = 8$ vertices in Fig. 2 has simple Laplacian spectrum and has $k = \frac{n}{2} - 1 = 3$ twins given by $\{v_1, v_2\}$, $\{v_3, v_4\}$, and $\{v_5, v_6\}$.

The proof of 3) is similar to 2) and is omitted. In this case, the bound $t = \frac{n-1}{2}$ is attained by the graph on $n = 11$ vertices in Fig. 2 which has simple Laplacian spectrum and has $k = \frac{n-1}{2} = 5$ twins given by $\{v_1, v_2\}$, $\{v_3, v_4\}$, $\{v_5, v_6\}$, $\{v_7, v_8\}$, and $\{v_9, v_{10}\}$. ■




Fig. 2. Sample of graphs with maximal twins and simple Laplacian eigenvalues.

In view of Lemma III.1, we make the following definition.

Definition III.1: A simple graph G is said to be a *twin graph* if $L = L(G)$ has simple spectrum and G contains the maximal number of twins $t = \lfloor \frac{n-1}{2} \rfloor$.

Let $w(n)$ be the number of twin graphs on n vertices. We have verified numerically that $w(n) = 0$ for $n \leq 6$ and that $w(7) = 12$, $w(8) = 36$, $w(9) = 42$, and $w(10) = 924$. We conjecture that twin graphs exist for all $n \geq 7$.

If G is a twin graph with vertex set $V = \{v_1, v_2, \dots, v_n\}$, and n is even, we assume that the $t = \frac{n}{2} - 1$ twins are $C_1 = \{v_1, v_2\}, C_2 = \{v_3, v_4\}, \dots, C_t = \{v_{n-3}, v_{n-2}\}$ and that v_{n-1} and v_n are the nontwin vertices. In this case, we call $\pi^* = \{C_1, C_2, \dots, C_t, \{v_{n-1}\}, \{v_n\}\}$ the *twin partition* of G . If n is odd, then we may assume that the $t = \frac{n-1}{2}$ twins are $C_1 = \{v_1, v_2\}, C_2 = \{v_3, v_4\}, \dots, C_t = \{v_{n-2}, v_{n-1}\}$, and v_n will denote the nontwin vertex. In this case, $\pi^* = \{C_1, C_2, \dots, C_t, \{v_n\}\}$ is the twin partition. In either case, π^* is an equitable partition of G .

We end this section with a discussion on the *minimal controllability problem* for twin graphs [14]. Given a linear time-invariant system $\dot{x} = Mx$ on \mathbb{R}^n , the minimal controllability problem is to find a smallest subset $\mathcal{I} = \{i_1, \dots, i_p\} \subset \{1, 2, \dots, n\}$ such that if $B = [e_{i_1} \dots e_{i_p}] \in \mathbb{R}^{n \times p}$, then the linear time-invariant control system $\dot{x} = Mx + Bu$ is controllable. It was proved in [14] that finding such a smallest \mathcal{I} within a multiplicative factor of $c \log(n)$ is NP-hard for some absolute constant $c > 0$, even when M is symmetric. It is known that if the maximum geometric multiplicity of the eigenvalues of M is q , then $\text{rank}(B) \geq q$ whenever (M, B) is a controllable pair [20, pg. 95]. The existence of twin graphs shows that q is, in general, a very poor lower bound for $\text{rank}(B) = p$. Indeed, if G is a twin graph with twin cells C_1, C_2, \dots, C_t , then any subset \mathcal{I} chosen as input nodes such that $\mathcal{I} \cap C_i = \emptyset$ renders (L, B) uncontrollable; this follows easily since the Faria eigenvector associated to C_i is clearly orthogonal to every column of B . As a consequence, we obtain the following.

Theorem III.1: Let $G = (V, E)$ be a n -vertex twin graph with Laplacian matrix L and let $t = \lfloor \frac{n-1}{2} \rfloor$ denote the number of twins in G . Let $\mathcal{I} = \{i_1, \dots, i_p\} \subset \{1, 2, \dots, n\}$ and let $B = [e_{i_1} \dots e_{i_p}] \in \mathbb{R}^{n \times p}$. If (L, B) is controllable, then $p \geq t$.

The punchline of Theorem III.1 is that, at least for consensus-type dynamics, the generic case of simple eigenvalues in a graph matrix does not eliminate the necessity of controlling a significant fraction of the nodes to achieve network controllability.

IV. STRONGLY UNCONTROLLABLE GRAPHS

In this section, we present the main result of this article. For completeness, we formally state the definition of a strongly uncontrollable graph.

Definition IV.1: Let G be a simple connected graph and suppose that $L = L(G)$ has a simple spectrum. We say that G is *strongly uncontrollable* if the pair (L, b) is uncontrollable for every $b \in \{0, 1\}^n$.

Strong uncontrollability can also be defined using the adjacency or signless Laplacian matrix. However, the Laplacian matrix has the key property that it has the all ones vector e as an eigenvector with corresponding eigenvalue $\lambda = 0$ and, therefore, (L, b) is uncontrollable if and only if $(L, e - b)$ is uncontrollable (provided $n \geq 2$). Also, as will be seen below, our sufficient condition for strong uncontrollability relies heavily on the orthogonality of eigenvectors of L with the eigenvector e .

Although a twin graph G has many Faria eigenvectors, it is not necessarily a strongly uncontrollable graph. In fact, we have verified that all twin graphs on $n = 7$ vertices are not strongly uncontrollable and only 10 of the $w(8) = 36$ twin graphs on $n = 8$ vertices are strongly uncontrollable. Below, we give a sufficient condition for strong uncontrollability of twin graphs in terms of AEPs of the quotient graph G_{π^*} .

Theorem IV.1: Let G be a twin graph and suppose that $n = |V(G)|$ is even. Let $\pi^* = \{C_1, C_2, \dots, C_t, \{v_{n-1}\}, \{v_n\}\}$ denote the twin partition of G . If the quotient graph G_{π^*} has an AEP of the form $\rho = \{S_1, S_2, \dots, S_m, \{\{v_{n-1}\}\}, \{\{v_n\}\}\}$, then G is strongly uncontrollable.

Proof: By the PBH eigenvector controllability test, we must show that each binary vector $b \in \{0, 1\}^n$ is orthogonal to some eigenvector of L . If b is a binary vector such that $b_u = b_v$ where $\{u, v\} = C_i$ for some $i \in \{1, 2, \dots, t\}$, then b is orthogonal to the Faria eigenvector $e_u - e_v$, and, thus, a necessary condition for controllability is that $b_u \neq b_v$ for all twin pairs $\{u, v\}$. To prove the theorem, we will show the existence of two eigenvectors x and \tilde{x} of L that are orthogonal to the binary vectors that have exactly one nonzero entry on each twin cell of π^* . To that end, let P_{π^*} be the characteristic matrix of the twin partition π^* and let ξ_i be the characteristic vector of the cell C_i for $i = 1, 2, \dots, t$. Then, clearly

$$\text{range}(P_{\pi^*}) = \left\{ \alpha e_{n-1} + \beta e_n + \sum_{i=1}^t \gamma_i \xi_i : \alpha, \beta, \gamma_i \in \mathbb{R} \right\}.$$

Since $\{S_1, \dots, S_m\}$ is a partition of the set $\{C_1, C_2, \dots, C_t\}$, we can write for each $j \in \{1, 2, \dots, m\}$ that $S_j = \{C_{j,1}, C_{j,2}, \dots, C_{j,|S_j|}\}$, where $C_{j,i} \in \{C_1, C_2, \dots, C_t\}$ for all $i \in \{1, 2, \dots, |S_j|\}$. We may, therefore, write an arbitrary vector in $\text{range}(P_{\pi^*})$ in the form

$$\alpha e_{n-1} + \beta e_n + \sum_{j=1}^m \sum_{i=1}^{|S_j|} \gamma_{j,i} \xi_{j,i}$$

where $\xi_{j,i}$ is the characteristic vector of the cell $C_{j,i} \in S_j$, and $\alpha, \beta, \gamma_{j,i} \in \mathbb{R}$.

The partition $\rho = \{S_1, S_2, \dots, S_m, \{\{v_{n-1}\}\}, \{\{v_n\}\}\}$ is π^* -regular; indeed, each set S_j contains cells of π^* that have

cardinality two (twin cells) and the set $\{\{v_{n-1}\}, \{v_n\}\}$ clearly consists of cells of π^* of the same cardinality. Thus, by Lemma II.1, there exists an eigenvector y of $L(G_{\pi^*})$ such that $P_{\rho}^T y = 0$. Therefore, the eigenvector $x = P_{\pi^*} y$ of L can be written in the form

$$x = \alpha(e_{n-1} - e_n) + \sum_{j=1}^m \sum_{i=1}^{|S_j|} \gamma_{j,i} \xi_{j,i}$$

where for each $j \in \{1, 2, \dots, m\}$ we have

$$\sum_{i=1}^{|S_j|} \gamma_{j,i} = 0. \quad (1)$$

We note that there exists at least one $j \in \{1, 2, \dots, m\}$ such that $|S_j| \geq 2$ (i.e., at least one set S_j is not a singleton cell); if not then $\gamma_{j,i} = 0$ for all i, j and, therefore, $x = \alpha(e_{n-1} - e_n)$ which implies that $\{v_{n-1}, v_n\}$ is a twin cell of G which contradicts the maximality of t .

Since ρ is an AEP of G_{π^*} , there exists an eigenvector $\tilde{y} \neq e$ of $L(G_{\pi^*})$ such that $\tilde{y} \in \text{range}(P_{\rho})$. Hence, the eigenvector $\tilde{x} = P_{\pi^*} \tilde{y}$ of L takes the form

$$\tilde{x} = \beta(e_{n-1} + e_n) + \sum_{j=1}^m \mu_j \sum_{i=1}^{|S_j|} \xi_{j,i}$$

for $\mu_j \in \mathbb{R}$. Since \tilde{x} is orthogonal to the all ones eigenvector of L , and since each vector $\xi_{j,i}$ is a sum of two distinct standard basis vectors, we have

$$0 = \langle \tilde{x}, e \rangle = 2\beta + \sum_{j=1}^m 2|S_j| \mu_j$$

which simplifies to

$$\beta + \sum_{j=1}^m |S_j| \mu_j = 0. \quad (2)$$

Now let $b \in \{0, 1\}^n$. As already mentioned at the beginning of the proof, we need only consider the case that b has exactly one nonzero entry on each twin cell. Hence, $\langle b, \xi_{j,i} \rangle = 1$ for all $i \in \{1, \dots, |S_j|\}$ and all $j \in \{1, \dots, m\}$. There are three cases to consider. If $b_{n-1} = b_n = 1$, then from (1), we have

$$\langle x, b \rangle = \alpha - \alpha + \sum_{j=1}^m \sum_{i=1}^{|S_j|} \gamma_{j,i} = 0.$$

If $b_{n-1} = b_n = 0$, then again from (1), we have

$$\langle x, b \rangle = \sum_{j=1}^m \sum_{i=1}^{|S_j|} \gamma_{j,i} = 0.$$

Finally, if $b_{n-1} \neq b_n$, then from (2), we have

$$\langle \tilde{x}, b \rangle = \beta + \sum_{j=1}^m \mu_j |S_j| = 0.$$

Thus, in any case, b is orthogonal to an eigenvector of L and, thus, (L, b) is uncontrollable. Since G has simple eigenvalues, by definition G is a strongly uncontrollable graph. ■

The case of an odd number of vertices is similar.




Fig. 3. G and G_{π^*} satisfying Theorem IV.2.

Theorem IV.2: Let G be a twin graph and suppose that $n = |V(G)|$ is odd. Let $\pi^* = \{C_1, C_2, \dots, C_t, \{v_n\}\}$ denote the twin partition of G . If G_{π^*} has a nontrivial AEP of the form $\rho = \{S_1, S_2, \dots, S_m, \{\{v_n\}\}\}$, then G is strongly uncontrollable.

We illustrate the previous results with an example.

Example IV.1: The twin graph G on $n = 11$ vertices shown in Fig. 3 satisfies the assumptions of Theorem IV.2; we also display the quotient graph G_{π^*} where $\pi^* = V(G_{\pi^*}) = \{C_1, C_2, C_3, C_4, C_5, \{v_{11}\}\}$ is the twin partition of G . The Faria eigenvector associated to the twin C_i is $x_i = e_{2i-1} - e_{2i}$ for $i = 1, 2, \dots, 5$. The adjacency matrix of G_{π^*} is

$$A_{\pi^*} = \begin{bmatrix} 0 & 2 & 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 2 & 2 & 2 & 0 & 0 \end{bmatrix}.$$

One can verify by inspection that $\rho = \{\{C_1, C_5\}, \{C_2, C_4\}, \{C_3\}, \{v_{11}\}\}$ is an AEP of G_{π^*} . The ρ -merge of π^* is $\pi_{\rho}^* = \{\{1, 2, 9, 10\}, \{3, 4, 7, 8\}, \{5, 6\}, \{11\}\}$ and the quotient graph of $G_{\pi_{\rho}^*}$ has adjacency matrix

$$A(G_{\pi_{\rho}^*}) = \begin{bmatrix} 0 & 2 & 0 & 0 \\ 2 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 4 & 2 & 0 \end{bmatrix}.$$

Theorems IV.1 and IV.2 are not necessary for strong uncontrollability as will be seen in the next section where we consider the stability of strong uncontrollability to vertex additions. Through numerical investigations, however, we have found that all strongly uncontrollable graphs up to $n = 12$ vertices have at least three twin pairs. For the adjacency matrix, twin vertices induce the eigenvalue $\lambda = 0$ or $\lambda = -1$ depending on whether the twin vertices are adjacent or not adjacent. Thus, any graph with three twin vertices has an adjacency matrix with at least one repeated eigenvalue. This observation leads us to conjecture that strongly uncontrollable graphs using the adjacency matrix A do not exist. As for the signless Laplacian matrix $Q = D + A$, we have not found any strongly uncontrollable graphs up to $n = 10$

vertices. Although our numerical investigations are for very small n , they suggest that strong uncontrollability is a property that may only exist in consensus-type network dynamics.

V. UNCONTROLLABILITY UNDER PERTURBATIONS

In this section, we analyze the stability of the strong uncontrollability property to vertex additions. Our results rely on the following well-known result concerning the Laplacian eigenvalues under the graph join operation. Given two simple graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ such that $V_1 \cap V_2 = \emptyset$, the *join* of G_1 and G_2 is the graph $G = G_1 \vee G_2$ with vertex set $V(G) = V_1 \cup V_2$ and edge set $E(G) = E_1 \cup E_2 \cup \{\{u, v\} \mid u \in V_1, v \in V_2\}$.

Theorem V.1 (see [22]): Let G_1 and let G_2 be simple graphs on disjoint sets of n_1 and n_2 vertices, respectively. Let L_1 be the Laplacian matrix of G_1 , with eigenvectors x_1, x_2, \dots, x_{n_1} and corresponding eigenvalues $0 = \alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_{n_1}$. Let L_2 be the Laplacian matrix of G_2 , with eigenvectors y_1, y_2, \dots, y_{n_2} and corresponding eigenvalues $0 = \beta_1 \leq \beta_2 \leq \dots \leq \beta_{n_2}$. Let L be the Laplacian matrix of the join graph $G = G_1 \vee G_2$. The following hold:

- 1) For all $i = 2, \dots, n_1$, $[x_i^T \ 0_{n_2}^T]^T$ is an eigenvector of L with eigenvalue $n_2 + \alpha_i$.
- 2) For all $j = 2, \dots, n_2$, $[0_{n_1}^T \ y_j^T]^T$ is an eigenvector of L with eigenvalue $n_1 + \beta_j$.
- 3) $[n_2 x_1^T \ -n_1 y_1^T]^T$ is an eigenvector of L with eigenvalue $n_1 + n_2$.

The following theorem describes how a strongly uncontrollable graph can be constructed from a lower-order strongly uncontrollable graph while preserving its automorphism group. Henceforth, we denote by S_n the symmetric group on $\{1, 2, \dots, n\}$, that is, the group of all permutations on $\{1, 2, \dots, n\}$.

Theorem V.2: Let G be a strongly uncontrollable graph on n vertices. Let \tilde{G} be the graph on $n + 1$ vertices obtained from G by adding a vertex and connecting it to all the vertices of G . Then, \tilde{G} is a strongly uncontrollable graph if and only if the spectral radius of L is less than n . In this case, $\text{Aut}(G)$ and $\text{Aut}(\tilde{G})$ are equal when viewed as subgroups of the symmetric group S_{n+1} .

Proof: Let x_1, x_2, \dots, x_n be the eigenvectors of $L = L(G)$ with corresponding eigenvalues $0 = \lambda_1 < \lambda_2 < \dots < \lambda_n$. Applying Theorem V.1 to G and the graph with one vertex, the set $\{e, [x_2^T \ 0]^T, \dots, [x_n^T \ 0]^T, [e^T \ -n]^T\}$ consists of mutually orthogonal eigenvectors of $\tilde{L} = L(\tilde{G})$ with eigenvalues $0 = \lambda_1 < \lambda_2 + 1 < \dots < \lambda_n + 1 \leq n + 1$. If $\lambda_n < n$, then $n + 1$ is a simple eigenvalue of \tilde{L} . Consequently, the eigenvalues of \tilde{L} are distinct.

Now, if $b \in \{0, 1\}^{n+1}$, then since G is strongly uncontrollable, there exists an eigenvector $x_i \neq x_1$ of L such that $[x_i^T \ 0]^T$ is orthogonal to b . This proves that (\tilde{L}, b) is uncontrollable for every $b \in \{0, 1\}^{n+1}$.

To prove the second claim, since the spectral radius of L is less than n , no vertex of G has degree $n - 1$. Hence, the vertex v_{n+1} that was added to G to form \tilde{G} is the only vertex of \tilde{G}




Fig. 4. Strongly uncontrollable graph on $n = 8$ vertices and its induced strongly uncontrollable graph on $n = 9$ vertices using Theorem V.2.

with degree n . It follows that any automorphism of \tilde{G} must fix v_{n+1} , and this proves that $\text{Aut}(G)$ and $\text{Aut}(\tilde{G})$ are equal when viewed as subgroups of S_{n+1} .

To prove the converse statement, assume that \tilde{G} is strongly uncontrollable. Then, by definition \tilde{L} has simple spectrum. Then, since $n + 1$ is an eigenvalue of \tilde{L} it follows that $\lambda_n < n$. ■

Theorem V.2 shows that if G is a twin graph on an even number of vertices, then \tilde{G} is not a twin graph, and, thus, showing that the property of being a twin graph is not necessary for strong uncontrollability.

Example V.1: Let $s(n)$ be the number of strongly uncontrollable graphs on n vertices. We have numerically verified that $s(8) = 10$ and $s(9) = 12$. All ten strongly uncontrollable graphs for $n = 8$ have spectral radius less than n and are all twin graphs. Hence, each strongly uncontrollable graph on $n = 8$ vertices induces a strongly uncontrollable graph on $n = 9$ vertices via Theorem V.2, none of which is a twin graph. One such pair is displayed in Fig. 4. The other two uncontrollable graphs on $n = 9$ vertices are twin graphs.

We now consider the case of adding two vertices. If $\tau : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ is a permutation such that $\tau(i) = j$ and $\tau(j) = i$ (with $i \neq j$) and fixes all other integers (i.e., τ is a transposition), we denote τ by $\tau = (i \ j)$.

Theorem V.3: Let G be a strongly uncontrollable graph with vertex set $V = \{v_1, v_2, \dots, v_n\}$. Let $G_2 = (\{v_{n+1}, v_{n+2}\}, \emptyset)$ be the empty graph on two vertices and let $\tilde{G} = G \vee G_2$. Then, \tilde{G} is a strongly uncontrollable graph if and only if $n - 2$ and n are not eigenvalues of L . In this case, $\text{Aut}(\tilde{G})$ is generated by the union of a generating set of $\text{Aut}(G)$ and the transposition $\tau = (n + 1 \ n + 2)$.

Proof: Let x_1, x_2, \dots, x_n be the eigenvectors of $L = L(G)$ with corresponding eigenvalues $0 = \lambda_1 < \lambda_2 < \dots < \lambda_n$. The Laplacian matrix of \tilde{G} can be written as

$$\tilde{L} = \begin{bmatrix} (L + 2I) & -e & -e \\ -e^T & n & 0 \\ -e^T & 0 & n \end{bmatrix}.$$

The vectors $e, [x_2^T \ 0]^T, \dots, [x_n^T \ 0]^T$ are eigenvectors of \tilde{L} with eigenvalues $0 = \lambda_1 < \lambda_2 + 2 < \dots < \lambda_n + 2$, and $[2e^T \ -n \ -n]^T$ is an eigenvector of \tilde{L} with eigenvalue $n + 2$. The Faria eigenvector $[0^T \ 1 \ -1]^T$ of \tilde{L} has corresponding

eigenvalue n . Thus, if $n - 2$ and n are not eigenvalues of L , then \tilde{L} has simple spectrum. Since G is a strongly uncontrollable graph, every binary vector in $\{0, 1\}^{n+2}$ is orthogonal to some eigenvector $[x_i^T \ 0 \ 0]^T$ of \tilde{L} . Thus, \tilde{G} is also a strongly uncontrollable graph. As in Theorem V.2, the converse statement is straightforward.

To prove the second statement, it is clear that every automorphism of G can be extended to an automorphism of \tilde{G} by asking that it fix the vertices v_{n+1} and v_{n+2} . The transposition $\tau = (n+1 \ n+2)$ is an automorphism of \tilde{G} and there are no other automorphisms of \tilde{G} that do not fix v_{n+1} and v_{n+2} . ■

Remark V.1: The procedure in Theorems V.2 and V.3 of taking a strongly uncontrollable graph G and creating a new strongly uncontrollable graph by joining it to an empty graph with n_2 vertices cannot be extended to the case $n_2 > 2$. Indeed, if $n_2 > 2$ vertices are joined to G , then \tilde{G} will contain an automorphism of order greater than two and, thus, \tilde{G} would not have distinct eigenvalues.

The following theorem can be seen as a complement of Theorem V.2.

Theorem V.4: Let G be a strongly uncontrollable graph with vertex set $V = \{v_1, v_2, \dots, v_n\}$ and suppose that $\deg(v_n) = n - 1$. Let \tilde{G} be the graph on $n + 1$ vertices obtained from G by adding a vertex and connecting it only to v_n . Then, \tilde{G} is a strongly uncontrollable graph if and only if 1 is not an eigenvalue of L . Moreover, $\text{Aut}(G)$ and $\text{Aut}(\tilde{G})$ are equal when viewed as subgroups of S_{n+1} .

Proof: Let x_1, x_2, \dots, x_n be eigenvectors of $L = L(G)$ with corresponding eigenvalues $0 = \lambda_1 < \lambda_2 < \dots < \lambda_n = n$. Since $\deg(v_n) = n - 1$, we may take $x_n = -e + ne_n$. By orthogonality of eigenvectors of L , we have $0 = \langle x_j, x_n \rangle = n \langle x_j, e_n \rangle$ for $2 \leq j \leq n - 1$, that is, $\langle x_j, e_n \rangle = e_n^T x_j = 0$. Now, $\tilde{L} = L(\tilde{G})$ takes the form

$$\tilde{L} = \begin{bmatrix} L + e_n e_n^T & -e_n \\ -e_n^T & 1 \end{bmatrix}$$

and, therefore, if we set $\tilde{x}_j = [x_j^T \ 0]^T$, for $2 \leq j \leq n - 1$, we have

$$\tilde{L}\tilde{x}_j = \begin{bmatrix} Lx_j + e_n e_n^T x_j \\ -e_n^T x_j \end{bmatrix} = \begin{bmatrix} Lx_j \\ 0 \end{bmatrix} = \lambda_j \tilde{x}_j.$$

Hence, \tilde{x}_j is an eigenvector of \tilde{L} with eigenvalue λ_j , for $2 \leq j \leq n - 1$. Now, $\tilde{x}_{n+1} = \tilde{L}e_n = [-1 \ -1 \ \dots \ -1 \ n \ -1]^T \in \mathbb{R}^{n+1}$ is an eigenvector of \tilde{L} with eigenvalue $\lambda_{n+1} = n + 1$. Finally, consider the vector $\tilde{x}_n = [-1 \ -1 \ \dots \ -1 \ 0 \ n - 1]^T \in \mathbb{R}^{n+1}$. A straightforward calculation shows that \tilde{x}_n is an eigenvector of \tilde{L} with eigenvalue 1. Hence, if $\lambda_j \neq 1$ for $2 \leq j \leq n - 1$, then \tilde{L} has simple eigenvalues $\{0, 1, \lambda_2, \lambda_3, \dots, \lambda_{n-1}, n + 1\}$. The rest of the proof is similar to that of Theorem V.2 and is omitted. ■

VI. CONCLUSION

In this article, we have characterized network topologies under which the Laplacian consensus dynamics are uncontrollable

for any subset of the nodes chosen as control inputs and that emit a common control signal. In these network topologies, the lack of controllability is not due to repeated eigenvalues of the Laplacian matrix, but instead is characterized by the structural properties of the network, namely, the existence of a maximal number of twin nodes and certain AEPs. We provided a sufficient condition for a network to contain this strong uncontrollability property and described network perturbations that leave the uncontrollability property invariant. We also related our work with the minimal controllability problem and showed how these network topologies require the control of essentially half of the nodes for any chance of controllability.

APPENDIX A PROOF OF LEMMA II.1

Before we give the proof of Lemma II.1, we need some preliminary results. Let $\pi = \{C_1, C_2, \dots, C_k\}$ be a partition of $V = \{v_1, \dots, v_n\}$ and let $K = \text{diag}(|C_1|, |C_2|, \dots, |C_k|) \in \mathbb{R}^{k \times k}$. It is easy to see that $K = P_\pi^T P_\pi$. Let $\rho = \{S_1, S_2, \dots, S_m\}$ be a partition of π and let $\xi_j \in \{0, 1\}^k$ be the characteristic vector of S_j . If ρ is π -regular (i.e., all cells in S_j have the same cardinality), then clearly

$$K\xi_j = |C_i|\xi_j$$

for any (and hence all) $C_i \in S_j$. It follows then that $KP_\rho = P_\rho \tilde{K}$ where $\tilde{K} = \text{diag}(|C_{1,1}|, |C_{2,1}|, \dots, |C_{m,1}|)$ where $C_{j,1} \in S_j$ for $j = 1, 2, \dots, m$. An identical argument shows that $K^{-1}P_\rho = P_\rho \tilde{K}^{-1}$. We can now prove Lemma II.1.

Proof of Lemma II.1: Let $\pi = \{C_1, C_2, \dots, C_k\}$ and let $K = P_\pi^T P_\pi = \text{diag}(|C_1|, |C_2|, \dots, |C_k|)$ as above. Then, from Theorem II.1, we have that $L_\pi = K^{-1}P_\pi^T L P_\pi$. Also, from Theorem II.1 applied to the quotient graph G_π and the partition ρ , we have that $L_\pi P_\rho = P_\rho L_{\pi_\rho}$, where π_ρ is the ρ -merge of π . Then

$$\begin{aligned} L_\pi^T P_\rho &= (P_\pi^T L^T P_\pi K^{-1}) P_\rho \\ &= (P_\pi^T L P_\pi)(K^{-1} P_\rho) \\ &= (K L_\pi)(K^{-1} P_\rho) \\ &= (K L_\pi)(P_\rho \tilde{K}^{-1}) \\ &= K P_\rho L_{\pi_\rho} \tilde{K}^{-1} \\ &= P_\rho (\tilde{K} L_{\pi_\rho} \tilde{K}^{-1}). \end{aligned}$$

In other words, $P_\rho^T L_\pi = (\tilde{K} L_{\pi_\rho} \tilde{K}^{-1})^T P_\rho^T$, and, thus, if $P_\rho^T x = 0$, then clearly $P_\rho^T L_\pi x = 0$, i.e., $\ker(P_\rho^T)$ is L_π -invariant. ■

ACKNOWLEDGMENT

The author would like to thank the anonymous referees for their comments and suggestions that improved the presentation of the paper.

REFERENCES

- [1] H. Tanner, "On the controllability of nearest neighbor interconnections," in *Proc. IEEE Conf. Decis. Control*, 2004, pp. 2467–2472.
- [2] A. Rahmani, M. Ji, M. Mesbahi, and M. Egerstedt, "Controllability of multi-agent systems from a graph-theoretic perspective," *SIAM J. Control Optim.*, vol. 48, no. 1, pp. 162–186, 2009.
- [3] S. Martini, M. Egerstedt, and A. Bicchi, "Controllability analysis of multi-agent systems using relaxed equitable partitions," *Int. J. Syst., Control Commun.*, vol. 2, no. 1, pp. 100–121, 2010.
- [4] G. Parlangeli and G. Notarstefano, "On the reachability and observability of path and cycle graphs," *IEEE Trans. Autom. Control*, vol. 57, no. 3, pp. 743–748, Mar. 2012.
- [5] A. Chapman, M. Nabi-Abdolyousefi, and M. Mesbahi, "Controllability and observability of network-of-networks via Cartesian products," *IEEE Trans. Autom. Control*, vol. 59, no. 10, pp. 2668–2679, Oct. 2014.
- [6] S. Zhang, M. Cao, and M. Camlibel, "Upper and lower bounds for controllable subspaces of networks of diffusively coupled agents," *IEEE Trans. Autom. Control*, vol. 59, no. 3, pp. 745–750, Mar. 2014.
- [7] N. Monshizadeh, S. Zhang, and M. Camlibel, "Zero forcing sets and controllability of dynamical systems defined on graphs," *IEEE Trans. Autom. Control*, vol. 59, no. 9, pp. 2562–2567, Sep. 2014.
- [8] A. Chapman and M. Mesbahi, "On symmetry and controllability of multi-agent systems," in *Proc. IEEE 53rd Annu. Conf. Decis. Control*, 2014, pp. 625–630.
- [9] C. Aguilar and B. Ghahesifard, "Graph controllability classes for the Laplacian leader-follower dynamics," *IEEE Trans. Autom. Control*, vol. 60, no. 6, pp. 1611–1623, Jun. 2015.
- [10] C. Aguilar and B. Ghahesifard, "Almost equitable partitions and new necessary conditions for network controllability," *Automatica*, vol. 80, pp. 25–31, 2017.
- [11] A. Yazıcıoğlu, W. Abbas, and M. Egerstedt, "Graph distances and controllability of networks," *IEEE Trans. Autom. Control*, vol. 61, no. 12, pp. 4125–4130, Dec. 2016.
- [12] S.-P. Hsu, "Constructing a controllable graph under edge constraints," *Syst. Control Lett.*, vol. 107, pp. 110–116, 2017.
- [13] Y. Guan and L. Wang, "Controllability of multi-agent systems with directed and weighted signed networks," *Syst. Control Lett.*, vol. 116, pp. 47–55, 2018.
- [14] A. Olshevsky, "Minimal controllability problems," *IEEE Trans. Control Netw. Syst.*, vol. 1, no. 3, pp. 249–258, Sep. 2014.
- [15] A. Olshevsky, "Minimum input selection for structural controllability," in *Proc. Amer. Control Conf.*, 2015, pp. 2218–2223.
- [16] S. Pequito, S. Kar, and P. Aguiar, "On the complexity of the constrained input selection problem for structural linear systems," *Automatica*, vol. 62, pp. 193–199, 2015.
- [17] S. Pequito, S. Kar, and P. Aguiar, "A framework for structural input/output and control configuration selection in large-scale systems," *IEEE Trans. Autom. Control*, vol. 61, no. 2, pp. 303–318, Feb. 2016.
- [18] S. Pequito, G. Ramos, S. Kar, P. Aguiar, and J. Ramos, "The robust minimal controllability problem," *Automatica*, vol. 82, pp. 261–268, 2017.
- [19] C. Commault and J.-M. Dion, "Input addition and leader selection for the controllability of graph-based systems," *Automatica*, vol. 49, no. 11, pp. 3322–3328, 2013.
- [20] E. Sontag, *Mathematical Control Theory: Deterministic Finite Dimensional Systems*, 2nd ed. Berlin, Germany: Springer, 1998.
- [21] B. MacArthur, R. Sánchez-García, and J. Anderson, "Symmetry in complex networks," *Discrete Appl. Math.*, vol. 156, no. 18, pp. 3525–3531, 2008.
- [22] T. Bıyıkoglu, J. Leydold, and P. Stadler, *Laplacian Eigenvectors of Graphs: Perron-Frobenius and Faber-Krahn Type Theorems*, vol. 1915. New York, NY, USA: Springer, 2007.
- [23] D. Cardoso, C. Delorme, and P. Rama, "Laplacian eigenvectors and eigenvalues and almost equitable partitions," *Eur. J. Combinatorics*, vol. 28, no. 3, pp. 665–673, 2007.
- [24] N. Monshizadeh, H. Trentelman, and M. Camlibel, "Projection-based model reduction of multi-agent systems using graph partitions," *IEEE Trans. Control Netw. Syst.*, vol. 1, no. 2, pp. 145–154, Jun. 2014.
- [25] N. Monshizadeh, S. Zhang, and M. Camlibel, "Disturbance decoupling problem for multi-agent systems: A graph topological approach," *Syst. Control Lett.*, vol. 76, pp. 35–41, 2015.
- [26] I. Stewart, M. Golubitsky, and M. Pivato, "Symmetry groupoids and patterns of synchrony in coupled cell networks," *SIAM J. Appl. Dynamical Syst.*, vol. 2, no. 4, pp. 609–646, 2003.
- [27] M. Golubitsky, I. Stewart, and A. Török, "Patterns of synchrony in coupled cell networks with multiple arrows," *SIAM J. Appl. Dynamical Syst.*, vol. 4, no. 1, pp. 78–100, 2005.
- [28] D. Cvetković, P. Rowlinson, and S. Simic, *Eigenspaces of Graphs* Issue 66 of Encyclopedia of Mathematics and Its Applications. New York, NY, USA: Cambridge Univ. Press, 1997.



Cesar O. Aguilar received the B.Sc. degree in mathematics and engineering from Queen's University, Kingston, ON, Canada, in 2003 and the M.Sc. degree in electrical and computer engineering from the University of Alberta, Edmonton, AB, Canada, in 2005. He received the Ph.D. degree in mathematics from Queen's University in 2010.

He held a National Research Council Postdoctoral Fellowship with the Department of Applied Mathematics, Naval Postgraduate School, Monterey, CA, USA, from 2010 to 2013. From 2013 to 2016, he was an Assistant Professor with the Department of Mathematics, California State University, Bakersfield, CA, USA. Since 2016, he has been with the Department of Mathematics, State University of New York at Geneseo, Geneseo, NY, USA. His research interests include nonlinear controllability theory, output regulation, optimal control, numerical methods in control theory, networked dynamical systems, and spectral graph theory.